

# Inequalities for Continued Fractions, II

Zaizhao Meng

School of Mathematical Sciences  
Peking University  
Beijing, 100871, P.R.China  
E-mail: mzzh@math.pku.edu.cn

In this paper, we investigate the monotone property of the continued fractions  $G(m, \lambda)$  as a function of  $m$  and  $\lambda$ . In particular, we obtain new inequality for the relative continued fractions.

**Key Words:** continued fractions; monotone property.

## 1. Introduction

We use the symbols and the properties of continued fractions in [1].

Let

$$\begin{aligned} \langle x_0, x_1, x_2, \dots \rangle &= x_0 + \frac{1}{x_1 + \frac{1}{x_2 + \frac{1}{\ddots}}}, \\ \langle x_0, x_1, x_2, \dots, x_n \rangle &= x_0 + \frac{1}{x_1 + \frac{1}{x_2 + \frac{1}{\ddots + \frac{1}{x_n}}}}, \end{aligned}$$

and  $P_{-2} = 0$ ,  $P_{-1} = 1$ ,  $Q_{-2} = 1$ ,  $Q_{-1} = 0$ ,  $P_n = x_n P_{n-1} + P_{n-2}$ ,

$Q_n = x_n Q_{n-1} + Q_{n-2}$ , then, for  $n \geq 0$ , we have

$$\langle x_0, x_1, x_2, \dots, x_n \rangle = \frac{P_n}{Q_n}.$$

Suppose that  $m > -1$ ,  $\lambda > 0$ , and let  $x_0 = m\lambda$ ,  $x_1 = (m+1)\lambda$ ,  $\dots$ ,  $x_j = (m+j)\lambda$ ,  $\dots$ ,  $x_n = (m+n)\lambda$ . We define the functions

$$G_n(m, \lambda) := \frac{P_n}{Q_n},$$

$$G(m, \lambda) := \lim_{n \rightarrow \infty} G_n(m, \lambda).$$

The purpose of this paper is to obtain new inequality for  $G(m, \lambda)$ .

**THEOREM .** For  $m \geq 0$ ,  $\lambda > 0$ , we have

$$G(m+1, \lambda) > \frac{m\lambda}{2} + \sqrt{\frac{m^2\lambda^2}{4} + 1} > G(m, \lambda).$$

In particular, for  $m \geq 1$ ,  $\lambda > 0$ ,  $G(m, \lambda) > 1$ .

## 2. Proof of Theorem

By Corollary 2 of [1], for  $\lambda > 0$ , we have

$$G(m+1, \lambda) > G(m, \lambda) = m\lambda + \frac{1}{G(m+1, \lambda)},$$

consequently, we obtain  $G(m+1, \lambda) > \frac{m\lambda}{2} + \sqrt{\frac{m^2\lambda^2}{4} + 1} > G(m, \lambda)$ , the theorem follows.

We obtain  $G(0, \lambda) = \frac{1}{G(1, \lambda)} < 1$ , for fixed  $\lambda > 0$ , there exists  $\alpha(\lambda) \in (0, 1)$  such that  $G(\alpha(\lambda), \lambda) = 1$ , by Theorem 3 of [1], for  $m > -1$ ,  $\lim_{\lambda \rightarrow 0+} G(m, \lambda) = 1$ , hence, there exists  $m \in (0, 1)$ , the function  $G(m, \lambda)$  of  $\lambda$  is not monotonically increasing in  $(0, +\infty)$ .

## References

- [1] Z.Z.Meng, *Inequalities for Continued Fractions*, Adv.in Math.(China) 35(2)(2006),143-154.